


**Amendments To The Claims:**

This listing of claims will replace all prior versions, and listings, of claims in the application:

What is claimed is:

1. – 25. (Canceled)

26. (Original) A compound



including salts, solvates, and pharmacologically functional derivatives thereof.

27. (Canceled)

28. (Currently Amended) A pharmaceutical composition comprising a compound according to claims ~~1 to~~ 26, and a pharmaceutically acceptable carrier.

29. (Currently Amended) A compound according to claims ~~1 to~~ 26 for use as an active therapeutic substance.

30. (Currently Amended) A compound according to claims ~~1 to~~ 26 for use in the treatment or prophylaxis of conditions or disorders affected by selective estrogen receptor modulation.

31. (Original) The compound of claim 30 wherein treatment or prophylaxis relates to osteoporosis, bone demineralization, reduced bone mass, density, or growth, osteoarthritis, acceleration of bone fracture repair and healing, acceleration of healing in joint replacement, periodontal disease, acceleration of tooth repair or growth, Paget's disease, osteochondrodysplasias, muscle wasting, the maintenance and enhancement of muscle strength and function, frailty or age-related functional decline ("ARFD"), sarcopenia, chronic fatigue syndrome, chronic myalgia, acute fatigue syndrome, acceleration of wound healing, maintenance of sensory function, chronic liver disease, AIDS,

weightlessness, burn and trauma recovery, thrombocytopenia, short bowel syndrome, irritable bowel syndrome, inflammatory bowel disease, Crohn's disease and ulcerative colitis, obesity, eating disorders including anorexia associated with cachexia or aging, hypercortisolism and Cushing's syndrome, cardiovascular disease or cardiac dysfunction, congestive heart failure, high blood pressure, breast cancer, malignant tumor cells containing the androgen receptor including breast, brain, skin, ovary, bladder, lymphatic, liver, kidney, uterine, pancreas, endometrium, lung, colon, and prostate, prostatic hyperplasia, hirsutism, acne, seborrhea, androgenic alopecia, anemia, hyperpilosity, adenomas and neoplasia of the prostate, hyperinsulinemia, insulin resistance, diabetes, syndrome X, dyslipidemia, urinary incontinence, atherosclerosis, libido enhancement, sexual dysfunction, depression, depressive symptoms, nervousness, irritability, stress, reduced mental energy and low self-esteem, improvement of cognitive function, endometriosis, polycystic ovary syndrome, counteracting preeclampsia, premenstrual syndrome, contraception, uterine fibroid disease, and/or aortic smooth muscle cell proliferation, vaginal dryness, pruritis, dyspareunia, dysuria, frequent urination, urinary tract infections, hypercholesterolemia, hyperlipidemia, peripheral vascular disease, restenosis, vasospasm, vascular wall damage due to immune responses, Alzheimer's disease, bone disease, aging, inflammation, rheumatoid arthritis, respiratory disease, emphysema, reperfusion injury, viral hepatitis, tuberculosis, psoriasis, systemic lupus erythematosus, amyotrophic lateral sclerosis, stroke, CNS trauma, dementia, neurodegeneration, breast pain and dysmenorrhea, menopausal or postmenopausal disorders, vasomotor symptoms, urogenital or vulvar vaginal atrophy, atrophic vaginitis, female sexual dysfunction, for enhancing libido, for the treatment of hypoactive sexual disorder, sexual arousal disorder, for increasing the frequency and intensity of orgasms, vaginismus, osteopenia, endometriosis, BPH (benign prostatic hypertrophy), dysmenorrhea, autoimmune diseases, Hashimoto's thyroiditis, SLE (systemic lupus erythematosus), myasthenia gravis, or reperfusion damage of ischemic myocardium.

32. (Original) The compound of claim 31 wherein treatment or prophylaxis relates to menopausal or postmenopausal disorders, vasomotor symptoms, urogenital or vulvar vaginal atrophy, atrophic vaginitis, endometriosis, female sexual dysfunction, breast cancer, depressive symptoms, diabetes, bone demineralization, or osteoporosis.

33. (Currently Amended) Use of a compound according to claims ~~1 to~~ 26 in the manufacture of a medicament for use in the treatment or prophylaxis of conditions or disorders associated with selective estrogen receptor modulation.
34. (Currently Amended) Use of a compound according to ~~any one of~~ claims ~~1 to~~ 26 in the manufacture of a medicament for use in the treatment or prophylaxis of osteoporosis, bone demineralization, reduced bone mass, density, or growth, osteoarthritis, acceleration of bone fracture repair and healing, acceleration of healing in joint replacement, periodontal disease, acceleration of tooth repair or growth, Paget's disease, osteochondrodysplasias, muscle wasting, the maintenance and enhancement of muscle strength and function, frailty or age-related functional decline ("ARFD"), sarcopenia, chronic fatigue syndrome, chronic myalgia, acute fatigue syndrome, acceleration of wound healing, maintenance of sensory function, chronic liver disease, AIDS, weightlessness, burn and trauma recovery, thrombocytopenia, short bowel syndrome, irritable bowel syndrome, inflammatory bowel disease, Crohn's disease and ulcerative colitis, obesity, eating disorders including anorexia associated with cachexia or aging, hypercortisolism and Cushing's syndrome, cardiovascular disease or cardiac dysfunction, congestive heart failure, high blood pressure, breast cancer, malignant tumore cells containing the androgen receptor including breast, brain, skin, ovary, bladder, lymphatic, liver, kidney, uterine, pancreas, endometrium, lung, colon, and prostate, prostatic hyperplasia, hirsutism, acne, seborrhea, androgenic alopecia, anemia, hyperpilosity, adenomas and neoplasia of the prostate, hyperinsulinemia, insulin resistance, diabetes, syndrome X, dyslipidemia, urinary incontinence, arteriosclerosis, libido enhancement, sexual dysfunction, depression, depressive symptoms, nervousness, irritability, stress, reduced mental energy and low self-esteem, improvement of cognitive function, endometriosis, polycystic ovary syndrome, counteracting preeclampsia, premenstrual syndrome, contraception, uterine fibroid disease, and/or aortic smooth muscle cell proliferation, vaginal dryness, pruritis, dyspareunia, dysuria, frequent urination, urinary tract infections, hypercholesterolemia, hyperlipidemia, peripheral vascular disease, restenosis, vasospasm, vascular wall damage due to immune responses, Alzheimer's disease, bone disease, aging, inflammation, rheumatoid arthritis, respiratory disease, emphysema, reperfusion injury, viral hepatitis, tuberculosis, psoriasis, systemic lupus erythematosus, amyotrophic lateral sclerosis, stroke, CNS trauma, dementia, neurodegeneration, breast

pain and dysmenorrhea, menopausal or postmenopausal disorders, vasomotor symptoms, urogenital or vulvar vaginal atrophy, atrophic vaginitis, female sexual dysfunction, for enhancing libido, for the treatment of hypoactive sexual disorder, sexual arousal disorder, for increasing the frequency and intensity of orgasms, vaginismus, osteopenia, endometriosis, BPH (benign prostatic hypertrophy), dysmenorrhea, autoimmune diseases, Hashimoto's thyroiditis, SLE (systemic lupus erythematosus), myasthenia gravis, or reperfusion damage of ischemic myocardium.

35. (Original) Use of a compound as in claim 34 wherein the condition or disorder is menopausal or postmenopausal disorders, vasomotor symptoms, urogenital or vulvar vaginal atrophy, atrophic vaginitis, endometriosis, female sexual dysfunction, breast cancer, depressive symptoms, diabetes, bone demineralization, or osteoporosis.

36. (Currently Amended) A method for the treatment or prophylaxis of conditions or disorders associated with selective estrogen receptor modulation comprising the administration of a compound according to ~~any one of claims 1 to 26~~.

37. (Currently Amended) A method for the treatment or prophylaxis of osteoporosis, bone demineralization, reduced bone mass, density, or growth, osteoarthritis, acceleration of bone fracture repair and healing, acceleration of healing in joint replacement, periodontal disease, acceleration of tooth repair or growth, Paget's disease, osteochondrodysplasias, muscle wasting, the maintenance and enhancement of muscle strength and function, frailty or age-related functional decline ("ARFD"), sarcopenia, chronic fatigue syndrome, chronic myalgia, acute fatigue syndrome, acceleration of wound healing, maintenance of sensory function, chronic liver disease, AIDS, weightlessness, burn and trauma recovery, thrombocytopenia, short bowel syndrome, irritable bowel syndrome, inflammatory bowel disease, Crohn's disease and ulcerative colitis, obesity, eating disorders including anorexia associated with cachexia or aging, hypercortisolism and Cushing's syndrome, cardiovascular disease or cardiac dysfunction, congestive heart failure, high blood pressure, breast cancer, malignant tumor cells containing the androgen receptor including breast, brain, skin, ovary, bladder, lymphatic, liver, kidney, uterine, pancreas, endometrium, lung, colon, and prostate, prostatic hyperplasia, hirsutism, acne, seborrhea, androgenic alopecia, anemia, hyperpilosity, adenomas and neoplasia of the prostate, hyperinsulinemia, insulin resistance, diabetes,

syndrome X, dyslipidemia, urinary incontinence, arteriosclerosis, libido enhancement, sexual dysfunction, depression, depressive symptoms, nervousness, irritability, stress, reduced mental energy and low self-esteem, improvement of cognitive function, endometriosis, polycystic ovary syndrome, counteracting preeclampsia, premenstrual syndrome, contraception, uterine fibroid disease, and/or aortic smooth muscle cell proliferation, vaginal dryness, pruritis, dyspareunia, dysuria, frequent urination, urinary tract infections, hypercholesterolemia, hyperlipidemia, peripheral vascular disease, restenosis, vasospasm, vascular wall damage due to immune responses, Alzheimer's disease, bone disease, aging, inflammation, rheumatoid arthritis, respiratory disease, emphysema, reperfusion injury, viral hepatitis, tuberculosis, psoriasis, systemic lupus erythematosus, amyotrophic lateral sclerosis, stroke, CNS trauma, dementia, neurodegeneration, breast pain and dysmenorrhea, menopausal or postmenopausal disorders, vasomotor symptoms, urogenital or vulvar vaginal atrophy, atrophic vaginitis, female sexual dysfunction, for enhancing libido, for the treatment of hypoactive sexual disorder, sexual arousal disorder, for increasing the frequency and intensity of orgasms, vaginismus, osteopenia, endometriosis, BPH (benign prostatic hypertrophy), dysmenorrhea, autoimmune diseases, Hashimoto's thyroiditis, SLE (systemic lupus erythematosus), myasthenia gravis, or reperfusion damage of ischemic myocardium.comprising the administration of a compound according to ~~any one of claims 1 to 26.~~

38. (Original) The method of claim 37 wherein the condition or disorder is menopausal or postmenopausal disorders, vasomotor symptoms, urogenital or vulvar vaginal atrophy, atrophic vaginitis, endometriosis, female sexual dysfunction, breast cancer, depressive symptoms, diabetes, bone demineralization, or osteoporosis.
39. (New) The compound of claim 26.
40. (New) A salt of claim 26.
41. (New) A solvate of claim 26.
42. (New) 4-[(4-Hydroxyphenyl)(3,3,5,5-tetramethylcyclohexylidene)methyl]benzoic acid.